

The role of primary to specialist care referral guidelines in cost effective care

Speaker: Dr Gillian Leng, Deputy Chief Executive, NICE

Authors: Mary Docherty, Tarang Sharma, Peter Littlejohns, Sarah Garner, Bhash Naidoo and Moni Choudhury
Research and Development, NICE


Background



- The UK National Health Service is facing significant financial pressures: **£20 billion efficiency savings needed by 2015**
- The Department of Health has launched an initiative to encourage savings through improvements in quality, innovation, productivity and prevention (**QIPP**)
- The National Institute for Health and Clinical Excellence (**NICE**) is responding by identifying priority areas where targeted advice may help **reduce ineffective practice and improve the quality of patient care.**

Primary to specialist care referrals

- In the UK, General Practitioners act as **gatekeepers** to specialist care
- Referral to a specialist service is a crucial point in a patient's management
- A delay or failure to refer when indicated could compromise patient care whilst unnecessary referrals are costly and can impact on the care of others
- Significant **variations** in referral practice are known to exist both within and across GP practices in the UK
- Variation in practice highlights a **potential target for quality and improvement savings.**

NHS variation in healthcare






Inappropriate referrals

- Inappropriate referral to specialist care places a **large financial burden** on the NHS
- By following best practice recommendations **clinical outcomes** and **patient experience** could be improved and **inequalities in patient care reduced**
- The impact of guidance on reducing variation in referral rates from primary to specialist care depends on the **quality and nature of the referral advice recommendations**
- A decision to refer a patient depends on several factors:
 - **The needs and expectations of individual patients and their families**
 - **The knowledge and experience of the individual practitioner**
 - **And the range, type and level of services available locally.**

 National Institute for
Health and Clinical Excellence

NICE and referral advice

- In 2001 NICE issued first advice on **referral to specialist services** for 11 common conditions seen in primary care
- Advice based on best evidence and consensus best practice
- Primary aim to encourage local health communities to discuss referral issues
- Subsequent referral advice issued **within topic specific clinical guideline documents.**



 National Institute for
Health and Clinical Excellence

Improving access to NICE referral advice

- As part of the QIPP initiative NICE revisited its referral advice recommendations
- A pilot project was initiated to improve the **accessibility** and **uptake** of NICE referral recommendations
- All NICE referral advice available in **clinical guidelines, cancer service guidance** and **public health guidance** were extracted and collated in a database.

Construction of a database

- Each record contains:
 - **the 'referral advice' recommendation**
 - **the timescale in which the referral should take place**
 - **any additional relevant information from the guideline.**
- The database is updated on a monthly basis incorporating all new NICE guidance – **currently 552!**
- GPs can search the database according to clinical field and specific clinical topic.

NHS
National Institute for Health and Clinical Excellence

Skip to content | Vision impaired | Login | Glossary | Contact | Sitemap | Site help

Search: GO

NICE NHS Evidence

Home Find guidance **Putting guidance into practice** Get involved News and press About NICE

Home... Using guidance... Referral advice

Benefits of implementation

NICE implementation programme

Commissioning guides supporting clinical service redesign

Implementation tools

Help implement NICE guidance

Education

Measuring the use of NICE guidance

Shared learning implementing NICE guidance

'Do not do' recommendations

Referral advice

NICE referral advice

What are the NICE 'referral advice' recommendations?

During the process of guidance development NICE's independent advisory bodies often give advice on referral from primary care to secondary care. At a [workshop](#) held in late 2009, it was suggested that referral advice should be made more accessible.

The NICE 'referral advice' recommendations database was created, and is maintained, by NICE's Research and Development team. [Read further information about the team's work.](#)

What is in the NICE 'referral advice' recommendations database?

The NICE 'referral advice' recommendations database contains current primary-to-secondary referral advice from NICE clinical guidelines, cancer service guidance and public health guidance. The database replaces the NICE publication 'Referral Advice: A Guide to Appropriate Referral from Primary to Specialist Services' (December 2001).

Each record in the database contains the 'referral advice' recommendation and includes additional information including the 'referral advice' category, health topic, the guidance it comes from (with a link to the relevant paragraph in the guidance) and the other 'referral advice' recommendations from the same guidance.

[Search the NICE 'referral advice' recommendations database](#)

Explanation of 'referral advice' categories denoting timescale in which a referral should be made

Each NICE 'referral advice' recommendation has been categorised as follows:

'Referral advice' category	Explanation
Immediate	Within 24 hours
Urgent	Within 14 days
Timeframe not specified	Timeframe to be determined by the clinician with responsibility for the referral

This page was last updated: 22 December 2010

Accessibility | Cymraeg | Freedom of Information | Data Protection | Copyright | Disclaimer | NHS Choices

Copyright © 2011 National Institute for Health and Clinical Excellence. All rights reserved.

Categories of advice

Each NICE 'referral advice' recommendation has been categorised as follows:

'Referral advice' category	Explanation
Immediate	Within 24 hours
Urgent	Within 14 days
Timeframe not specified	Timeframe to be determined by the clinician with responsibility for the referral

Development of referral advice requires a range of methods:

- **A systematic review of the evidence base**
- **Identification and consideration of other evidence not amenable to systematic review**
- **Consensus based on best practice using field experts, stakeholder opinion and extensive consultation.**

'Good' referral advice must be precise but also flexible.

Implementation tools

Help implement NICE guidance

Education

Measuring the use of NICE guidance

Shared learning implementing NICE guidance

'Do not do' recommendations

Referral advice

You can search the NICE 'referral advice' recommendations database by:

- typing in a specific term in the 'referral advice' recommendations box
- selecting an appropriate topic
- selecting the type of source guidance.

To see the full list of all NICE 'referral advice' recommendations leave all the fields blank and click the 'Search' button.

Search 'referral advice' recommendations

Referral advice' recommendation:

'Referral advice' by topic: Please select one

'Referral advice' by sub-topic: Please select one

'Referral advice' by guidance type: Please select one

Search

Search results

Results 1-10 of 29

Guidance ID	NICE 'referral advice' recommendation	'Referral advice' category	<input type="checkbox"/> Select all for export
CG15	Children and young people with suspected type 1 diabetes should be offered immediate (same day) referral to a multidisciplinary paediatric diabetes care team that has the competencies needed to confirm diagnosis and to provide immediate care.	Immediate	<input type="checkbox"/>
CG15	Children and young people with type 1 diabetes and suspected anxiety and/or depression should be referred promptly to child mental health professionals.	Time frame not specified	<input type="checkbox"/>
CG15	Diabetes care teams should consider referring children and young people with type 1 diabetes who have frequent hypoglycaemia and/or recurrent seizures for assessment of cognitive function, particularly if these occur at a young age.	Time frame not specified	<input type="checkbox"/>
CG15	Eye surveillance for adults newly diagnosed with type 1 diabetes should be started from diagnosis.	Time frame not specified	<input type="checkbox"/>
CG15	For people with an ulcerated foot, arrange referral to a specialist diabetes foot care team incorporating specifically trained foot care specialists (usually state-registered podiatrists) within 1-2 days if there is no overt infection of the ulcer or surrounding tissues, or as an emergency if such infection is present. If peripheral vascular disease is detected, refer for early assessment by a specialist vascular team.	Immediate	<input type="checkbox"/>

Home... Using guidance... Referral advice... 'Referral advice' recommendation details

Benefits of implementation

NICE implementation programme

Commissioning guides supporting clinical service redesign

Implementation tools

Help implement NICE guidance

Education

Measuring the use of NICE guidance

Shared learning implementing NICE guidance

'Do not do' recommendations

Referral advice

'Referral advice' recommendation details

Recommendation details

Referral advice recommendation: For people with an ulcerated foot, arrange referral to a specialist diabetes foot care team incorporating specifically trained foot care specialists (usually state-registered podiatrists) within 1-2 days if there is no overt infection of the ulcer or surrounding tissues, or as an emergency if such infection is present. If peripheral vascular disease is detected, refer for early assessment by a specialist vascular team.

Recommendation ID: 77

Referral advice category: Immediate

Source guidance details

Guidance ID: CG15

Guidance: Type 1 diabetes

Issue date: July 2004

Guidance document: View paragraph in relevant document

Paragraph number: 1.11.3.7

Page number: 59

View all NICE referral advice recommendations from this guidance:

- Children and young people with suspected type 1 diabetes should be offered immediate (same day) referral to a multidisciplinary paediatric diabetes care team that has the competencies needed to confirm diagnosis and to provide immediate care.
- Children and young people with type 1 diabetes and suspected anxiety and/or depression should be referred promptly to child mental health professionals.
- Diabetes care teams should consider referring children and young people with type 1 diabetes who have frequent hypoglycaemia and/or recurrent seizures for assessment of cognitive function, particularly if these occur at a young age.
- Eye surveillance for adults newly diagnosed with type 1 diabetes should be started from diagnosis.
- When managing retinopathy, referral to an ophthalmologist.

Overview of results

- **The project took longer** than anticipated revealing the difficulties in formulating specific and appropriate referral advice by guideline groups
- **Inconsistencies** in classification and methodologies used to define referral criteria were found between guideline groups
- **Terminology** relating to referral varied, making identification and extraction of all relevant information time-consuming
- Despite these difficulties an intuitive, comprehensive and up-to-date **database of referral advice** was produced.

Web stats for database use

Period	No. of page views	Cumulative total of page views
April 2011	815	20447
May 2011	997	21444
June 2011	845	22289
July 2011	516	22805
2010-2011	19632	19632


Following the launch of the database in **2010** web stats and reports from the NICE field team evidence that the **resource is continuing to be accessed.**

Referral advice and saving money

Example of cost savings with implementation:

Referral recommendation:	Source:	Estimated savings:
Referral for arthroscopic lavage and debridement should not be offered as part of treatment of osteoarthritis, unless the person has knee osteoarthritis with a clear history of mechanical locking (not gelling, 'giving way' or X-ray evidence of loose bodies).	Osteoarthritis Costing Report: Implementing NICE guidance	Estimated savings with implementation £23.6 million

Referral recommendations will be incorporated into **NICE clinical pathways**.



NICE Pathways

Pathway information Implementation Guidance

Home | About NICE Pathways | Contact us | Help | Browse pathways

Familial breast cancer: secondary care management

Breast cancer

Woman is referred to secondary care

```

    graph TD
      Start[Woman is referred to secondary care] --> Q1{Is the woman likely to be at raised risk or more than raised risk?}
      Q1 -- No --> Q2{Is the woman assessed to be at risk?}
      Q1 -- Yes --> Q3{Is there at least one criterion present in the family history?}
      Q2 -- No --> R1[Offer referral back to primary care]
      Q2 -- Yes --> Q3
      Q3 -- No --> Q4{Are there any unusual cancers in the family? Or is there a very strong paternal history of breast cancer? Or is there Jewish ancestry?}
      Q3 -- Yes --> R2[Offer referral to a specialist genetics service]
      Q4 -- No --> R3[Manage in secondary care]
      Q4 -- Yes --> R4[Seek advice from tertiary care contact about levels of risk and appropriateness of referral]
      R4 --> Q5{Are surveillance criteria met?}
      Q5 -- No --> R3
      Q5 -- Yes --> R5[Mammography and/or MRI surveillance after appropriate discussion of risks and benefits]
  
```

Manage in secondary care

Offer appropriate information.

All women at all care levels should receive standard written information that includes:

- risk information about population level and family history levels of risk, including a definition of family history
- the message that, if their family history alters, their risk may alter
- breast awareness information
- lifestyle advice, including information about HRT, oral contraceptives, lifestyle (including diet, alcohol etc.), breastfeeding, family size and timing
- contact details of those providing support and information, including local and national support groups
- the message that women can bring a family member/friend to appointments
- details of any appropriate trials or studies that may be appropriate to consider taking part in.

In addition women being cared for in primary care should receive:

- advice to return to discuss any implications if family history changes or breast symptoms develop.

Women being referred to secondary care should receive:

- information on the risk assessment that may take place and advice on obtaining a comprehensive family history if required
- information about potential outcomes, depending on the outcome of the risk assessment (including referral back to primary care, management in secondary care or referral to a specialist genetics service) and what might happen at each level.

Women being referred back to primary care should receive:

- detailed information about why secondary or tertiary care is not required
- advice to return to primary care to discuss any implications

Key priorities for implementation

Source guidance

Pathway created: May 2011 Last updated: May 2011
Copyright © 2011 National Institute for Health and Clinical Excellence. All Rights Reserved.

NICE Pathways

Pathway information Implementation Guidance

Home | About NICE Pathways | Contact us | Help | Browse pathways

Familial breast cancer: secondary care management

Breast cancer

Woman is referred to secondary care

```

    graph TD
      Start[Woman is referred to secondary care] --> Q1{Is the woman likely to be at raised risk or more than raised risk?}
      Q1 -- No --> Q2{Is the woman assessed to be at risk?}
      Q1 -- Yes --> Q3{Is there at least one criterion present in the family history?}
      Q2 -- No --> R1[Offer referral back to primary care]
      Q2 -- Yes --> Q3
      Q3 -- No --> Q4{Are there any unusual cancers in the family? Or is there a very strong paternal history of breast cancer? Or is there Jewish ancestry?}
      Q3 -- Yes --> R2[Offer referral to a specialist genetics service]
      Q4 -- No --> R3[Manage in secondary care]
      Q4 -- Yes --> R4[Seek advice from tertiary care contact about levels of risk and appropriateness of referral]
      R4 --> Q5{Are surveillance criteria met?}
      Q5 -- No --> R3
      Q5 -- Yes --> R5[Mammography and/or MRI surveillance after appropriate discussion of risks and benefits]
  
```

Manage in secondary care

Offer appropriate information.

All women at all care levels should receive standard written information that includes:

- risk information about population level and family history levels of risk, including a definition of family history
- the message that, if their family history alters, their risk may alter
- breast awareness information
- lifestyle advice, including information about HRT, oral contraceptives, lifestyle (including diet, alcohol etc.), breastfeeding, family size and timing
- contact details of those providing support and information, including local and national support groups
- the message that women can bring a family member/friend to appointments
- details of any appropriate trials or studies that may be appropriate to consider taking part in.

In addition women being cared for in primary care should receive:

- advice to return to discuss any implications if family history changes or breast symptoms develop.

Women being referred to secondary care should receive:

- information on the risk assessment that may take place and advice on obtaining a comprehensive family history if required
- information about potential outcomes, depending on the outcome of the risk assessment (including referral back to primary care, management in secondary care or referral to a specialist genetics service) and what might happen at each level.

Women being referred back to primary care should receive:

- detailed information about why secondary or tertiary care is not required
- advice to return to primary care to discuss any implications


Key priorities for implementation

Source guidance

Pathway created: May 2011 Last updated: May 2011
Copyright © 2011 National Institute for Health and Clinical Excellence. All Rights Reserved.

Implications for guideline developers

- Clear referral guidelines can **reduce costs** associated with inappropriate referrals
- Guideline developers can improve the quality, efficacy and uptake of advice by **agreeing uniform methods and terminology** when drawing up referral recommendations
- Referral advice must be both **clear and specific whilst leaving flexibility** to allow for clinical judgement
- Advice should be constructed in a way that encourages local health communities to **discuss referral problems and develop local referral protocols.**



Acknowledgements and contacts

Research and Development, Clinical and Public Health
Directorate:

<http://www.nice.org.uk/aboutnice/howwework/researchanddevelopment/about.jsp>

- **Authors:**
 - Dr Mary Docherty, Clinical Advisor
 - Ms Tarang Sharma, Senior Analyst
 - Professor Peter Littlejohns, Executive Director
 - Dr Sarah Garner, Associate Director
 - Dr Bhash Naidoo, Associate Director
 - Miss Moni Choudhury, Analyst
- **Contact:**
 - Tarang Sharma, Tel: +44 (0) 161 219 3876;
 - Email: Tarang.Sharma@nice.org.uk